

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07C 213/00, A61K 31/135, 31/54</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/22426</b> <b>(43) International Publication Date:</b> 28 May 1998 (28.05.98)
<b>(21) International Application Number:</b> PCT/SE97/01926 <b>(22) International Filing Date:</b> 18 November 1997 (18.11.97) <b>(30) Priority Data:</b> 9604253-6      20 November 1996 (20.11.96)      SE <b>(71) Applicant (for all designated States except US):</b> ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PALMER, Sven [SE/SE]; Rosenlundsvägen 28, VI, S-151 30 Södertälje (SE). SIDEN- QVIST, Michael [SE/SE]; Ålstigen 46, S-151 39 Södertälje (SE). <b>(74) Agent:</b> ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> NEW MANUFACTURING PROCESS OF METOPROLOL		
<b>(57) Abstract</b>  A method for the manufacture of metoprolol wherein the process is performed in water as solvent.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## NEW MANUFACTURING PROCESS OF METOPROLOL

Field of the invention

5 The present invention relates to an improved method for the manufacture of metoprolol base 1-(isopropylamino)-3-[p-(2-methoxyethyl)-phenoxy]-2-propanol) via the route of reacting p-(2-methoxyethyl)-phenol (A) and epichlorohydrin (B) and then reacting the obtained 1-(2,3-epoxypropoxy)-4-(2-methoxyethyl)-benzene (AB) with isopropylamine (C). The crude metoprolol base is then purified.

10

Prior art

Chemical Abstracts, vol. 112 (1990) abstract no 197820 discloses the reaction of p-(2-methoxyethyl)-phenol and epichlorohydrin in the two phase system of water and organic  
15 solvent.


Swedish patents 354 851 and 368 004 disclose the reaction of p-(2-methoxyethyl)-phenol and epichlorohydrin where the epichlorohydrin is used not only as a building block in the reaction but also as solvent.

20

Disclosure of the invention

It has now been found that metoprolol can be prepared in a manner that is fast, environmentally sound and gives a good yield and high purity using reactants that are  
25 known per se. The difference from the prior art is that the new method uses no other solvents than water for the reaction of A and B. From an environmental as well as an occupational hazard point of view it is a great advantage to be able to replace a hazardous organic solvent with a non-noxious solvent such as water.

30 The method of the invention is illustrated by the reaction scheme below:



General example

p-(2-Methoxyethyl)phenol (A) and epichlorohydrin (B, 1.4-2.0 eqv.) are reacted in water, at least 1.5 kg, preferably about 2 kg of water per kg of phenol, during the addition of sodium (or potassium) hydroxide solution, (1.3-1.7 eqv.) to form 1-(2,3-epoxypropoxy)-4-(2-methoxyethyl)benzene; (p-methoxyethyl-epoxypropoxybenzene). The reaction is preferably performed at a temperature of 50 - 70 °C.

The two phases are separated, and the p-methoxyethylepoxy-propoxybenzene is isolated by distillation under reduced pressure. More particular, the excess of epichlorohydrin is evaporated, and the epoxide is distilled under reduced pressure to obtain a product with a purity of about 96-98%. If desired, before distilling the main fraction of the epoxide, a prefraction/forecut (2-8 %, preferably 4-6 %) thereof could be distilled. The isolation by distillation of the epoxide under reduced pressure is an important part of the process and essential for the quality of the end product.

The epoxide is reacted with isopropylamine preferably in isopropyl alcohol to form metoprolol base. The amount of isopropylamine in relation to epoxide is at least 1 equivalent, preferably 3-6 equivalents. The reaction mixture is then treated in order to eliminate the excess of isopropylamine.

Alternatively, the amination with isopropylamine is carried out in a pressurized system without isopropyl alcohol at  $70 \pm 10^\circ\text{C}$  at pressures of 2.8-3.2 kg / 275-315 kPa.

The resulting metoprolol is dissolved in toluene, isobutyl methyl ketone or butyl acetate and extracted with dilute hydrochloric acid or sulphuric acid, preferably at pH 4-6. The phases are separated and the chosen solvent with sodium or potassium hydroxide solution to adjust the pH to 11 - 13 is added to the aqueous phase. The two phases are separated, and the organic phase is evaporated in vacuo to an oily residue of metoprolol base which is dissolved in acetone. Purified metoprolol base is then obtained by conventional means.

Working example1-(2,3-epoxypropoxy)-4-(2-methoxyethyl) benzene

5 p-(2-Methoxyethyl)phenol (A, ~ 6,6 mol), epichlorohydrin (B, 1.45 eqv.) and water (~ 2 kg) were combined and the mixture heated to ~ 50°C. Sodium hydroxide solution (50%; 1.4 eqv.) was added during 3 hours and the temperature was elevated to reach approximately 60°C during the addition. Formation of the title compound occurred during this period.

10

The batch was stirred for another hour at approximately 60°C, then cooled to approximately 50°C and the phases were separated and the product washed with water.

The residue was distilled at  $\leq 190^{\circ}\text{C}$  and a pressure of  $\leq 20$  mm Hg and the distillate was  
15 collected. The yield of the title compound was 80% of theory and the purity was 98% according to GC analysis.

Metoprolol base

20 1-(2,3-epoxypropoxy)-4-(2-methoxyethyl)benzene (1 kg, 4.8 mol), isopropyl alcohol (~0.9 kg) and isopropylamine (0.8-1.7 kg, 3-6 eqv.) were mixed and reacted for 2-5 hours at reflux. Formation of metoprolol base occurred during this period.

The reaction mixture was then concentrated at atmospheric pressure until the inner  
25 temperature reached ~100°C. Water was added to the batch and then distilled off in vacuo until the inner temperature reached ~100°C to form a concentrate.

The resulting concentrate was diluted with isobutyl methyl ketone (~0.6 kg) and water (~2.2 kg), and concentrated sulphuric acid was added, to adjust the pH to 4-6.

30

After separation, isobutyl methyl ketone (~ 1 kg) was added to the water layer, and concentrated sodium hydroxide solution was added to adjust the pH to 13.

The organic layer was concentrated in vacuo at  $\leq 80^{\circ}\text{C}$ , until distillation ceased, and the  
5 concentrated batch was redissolved in acetone (~1.6 kg) and filtered, to yield metoprolol base solution. The assay of metoprolol base in the solution was determined by titration. Yield: ~1.2 kg metoprolol base (100 %) ~95 % of theory. The purity of the metoprolol base was 96 %.

CLAIMS

1. A method for the manufacture of metoprolol, characterized by reacting in a first step p-(2-methoxyethyl)phenol and epichlorohydrin in water as solvent and at a temperature of 50 to 70°C, evaporating the excess of epichlorohydrin and then distilling the obtained 1-(2,3-epoxypropoxy)-4-(2-methoxyethyl)-benzene under reduced pressure, and in a second step reacting the obtained 1-(2,3-epoxypropoxy)-4-(2-methoxyethyl)-benzene and isopropylamine in the presence of isopropyl alcohol to form metoprolol base.
2. A method according to claim 1 wherein in the first step is carried out in the presence of sodium hydroxide.
3. A method according to claim 1 wherein in the first step is carried out in the presence of potassium hydroxide.
4. A method according to any of the preceding claims wherein the resulting metoprolol base is purified by dissolving the metoprolol base in a solvent selected among toluene, isobutyl methyl ketone and butyl acetate and extracted with either hydrochloric or sulphuric acid solution.
5. Metoprolol as prepared by the process according to any of claims 1-4.
6. A method as claimed in any one of claims 1 to 4 wherein the resulting metoprolol is converted into metoprolol tartrate.
7. A method as claimed in any one of claims 1 to 4 wherein the resulting metoprolol is converted into metoprolol succinate.



8. A method for the manufacture of a pharmaceutical preparation wherein metoprolol is produced by the method as claimed in any one of claims 1 to 4 or 6 to 7 and the metoprolol is thereafter formulated with a pharmaceutically acceptable diluent or carrier.

5

9. A method as claimed in claim 8 wherein a diuretic is included in the ingredients for the pharmaceutical preparation.

10. A method as claimed in claim 9 wherein the diuretic is hydrochlorothiazide.

10

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01926

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07C 213/00, A61K 31/135, A61K 31/54

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CASONLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ES 2011584 A6 (ESTEVE QUIMICA, S.A.), 16 January 1990 (16.01.90) --	1-8
X	Chemical Abstracts, Volume 113, No 21, 19 November 1990 (19.11.90), (Columbus, Ohio, USA), page 677, THE ABSTRACT No 190900r, ES, 2011584 A,, (Oranias Olsina, Gloria et al) 16 January 1990 (16.01.90) --	1-8
X	Chemical Abstracts, Volume 119, No 19, 8 November 1993 (08.11.93), (Columbus, Ohio, USA), page 863, THE ABSTRACT No 203128w, PL, 158497 A,, (Zjawiony, Jordan et al) 30 Sept 1992 (30.09.92) --	1-8

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

- \* Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

5 March 1998

Date of mailing of the international search report

09-03-1998

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01926

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9216197 A1 (SEPRACOR INC.), 1 October 1992 (01.10.92), the claims  --	8-10
A	Chemical Abstracts, Volume 112, No 21, 21 May 1990 (21.05.90), (Columbus, Ohio, USA), page 658, THE ABSTRACT No 197820c, PL, 144036 A,, (Zjawiony, Jordanet et al) 30 April 1988 (30.04.88)  -- -----	1-10

# INTERNATIONAL SEARCH REPORT

Information on patent family members

03/02/98

International application No.

PCT/SE 97/01926

Patent document cited in search report			Publication date	Patent family member(s)	Publication date
ES	2011584	A6	16/01/90	NONE	
ES	2011584	A	16/01/90	NONE	
PL	158497	A	30/09/92	NONE	
WO	9216197	A1	01/10/92	AT 141049 T AU 660984 B AU 1774192 A CA 2106494 A DE 69212646 D,T EP 0576617 A,B SE 0576617 T3 EP 0677291 A ES 2093833 T JP 6505994 T US 5362757 A	15/08/96 13/07/95 21/10/92 19/09/92 13/03/97 05/01/94 18/10/95 01/01/97 07/07/94 08/11/94
PL	144036	A	30/04/88	NONE	